arXiv:1409.3396v3 [quant-ph] 12 Jan 2015

More about the doubling degeneracy operators associated with Majorana fermions and Yang-Baxter equation

Li-Wei Yu[1,*] and Mo-Lin Ge[1]

[1]*Theoretical Physics Division, Chern Institute of Mathematics, Nankai University, Tianjin 300071, China*

A new realization of doubling degeneracy based on emergent Majorana operator Γ presented by Lee-Wilczek has been made. The Hamiltonian can be obtained through the new type of solution of Yang-Baxter equation, i.e. $\breve{R}(\theta)$-matrix. For 2-body interaction, $\breve{R}(\theta)$ gives the "superconducting" chain that is the same as 1D Kitaev chain model. The 3-body Hamiltonian commuting with Γ is derived by 3-body $\breve{R}_{123}$-matrix, we thus show that the essence of the doubling degeneracy is due to $[\breve{R}(\theta), \Gamma] = 0$. We also show that the extended Γ'-operator is an invariant of braid group B_N for odd N. Moreover, with the extended Γ'-operator, we construct the high dimensional matrix representation of solution to Yang-Baxter equation and find its application in constructing $2N$-qubit Greenberger-Horne-Zeilinger state for odd N.

PACS numbers: 71.10.Pm, 74.20.Mn, 02.10.De, 03.67.Mn,

The Majorana mode [1–4] has attracted increasing attention in physics due to its potential applications in topological quantum information processing [5–7]. Specifically, the degenerate ground state in Majorana mode serves as topologically protected states which can be used for topological quantum memory.

In the Ref. [8], Lee and Wilczek presented a new operator Γ that provided the doubling degeneracy for the Hamiltonian formed by Majorana fermions to overcome the conceptional incompletion of the algebraic set for the Majorana model. Following the Ref. [8], the Majorana operators γ_i's satisfy Clifford algebraic relations:

$$\{\gamma_i, \gamma_j\} = 2\delta_{ij}, \tag{1}$$

and the Hamiltonian takes the form

$$H_{\text{int}} = -\mathrm{i}\left(\alpha\gamma_1\gamma_2 + \beta\gamma_2\gamma_3 + \kappa\gamma_3\gamma_1\right). \tag{2}$$

The algebra in equation (1) is conceptually incomplete. Besides the parity, the nonlinear operator Γ is introduced [8]

$$\Gamma = -\mathrm{i}\gamma_1\gamma_2\gamma_3 \tag{3}$$

to form the set

$$\begin{gathered}\Gamma^2 = 1, P^2 = 1, [\Gamma, H_{\text{int}}] = 0, [P, H_{\text{int}}] = 0, \\ [\Gamma, \gamma_j] = 0, \{P, \gamma_j\} = 0, \{\Gamma, P\} = 0,\end{gathered} \tag{4}$$

where P implements the electron number parity, and $P^2 = 1$. The emergent Majorana operator Γ and parity operator P lead to the doubling degeneracy at any energy level, not only for the ground state.

On the other hand, based on the obtained new type of solution $\breve{R}_i(\theta)$ of Yang-Baxter equation (YBE), which is related to Majorana operators, the corresponding Hamiltonian can be found by following the standard way [9], i.e. the Hamiltonian $H \sim \frac{\partial \breve{R}_i(\theta)}{\partial\theta}|_{\theta=0}$. We find that the Hamiltonian derived from $\breve{R}_i(\theta)$ is 1D Kitaev model [1]. Moreover, because 1+1D 3-body S-matrix can be decomposed into three 2-body S-matrices based on YBE, we construct the 3-body Hamiltonian from 3-body S-matrix and find its doubling degeneracy. Hence, the advantage of parametrizing the braiding operator B_i to $\breve{R}_i(\theta)$ is that the desired Hamiltonian associated with Majorana operators can be derived from $\breve{R}_i(\theta)$.

Now let us first give a brief introduction to the Majorana representation of braiding operator as well as the solution of Yang-Baxter equation.

The non-Abelian statistics [10] of Majorana fermion (MF) has been proposed in both 1D quantum wires network [7] and 2D $p + \mathrm{i}p$ superconductor [2].

*Electronic address: NKyulw@gmail.com

For $2N$ Majorana fermions, the braiding operators of Majorana fermions form braid group B_{2N} generated by elementary interchanges $B_i = U_{i,i+1} = \exp(\frac{\pi}{4}\gamma_i\gamma_{i+1})$ of neighbouring particles ($i = 1, 2\cdots 2N-1$) with the following braid relations:

$$B_iB_{i+1}B_i = B_{i+1}B_iB_{i+1}, \tag{5}$$
$$B_iB_j = B_jB_i, \quad |i-j| > 1. \tag{6}$$

The Yang-Baxter equation (YBE) [9, 11, 12] is a natural generalization of braiding relation with the parametrized form:

$$\breve{R}_i(x)\breve{R}_{i+1}(xy)\breve{R}_i(y) = \breve{R}_{i+1}(y)\breve{R}_i(xy)\breve{R}_{i+1}(x), \tag{7}$$

where x, y stand for spectral parameters,

$$\breve{R}_i = \frac{1}{\sqrt{1+x^2}}(B_i + xB_i^{-1}). \tag{8}$$

The solutions of equation (7) was intensively studied by Yang, Baxter, Faddeev and other authors [11–20] in dealing with many body problems, statistical models, low-dimensional quantum field theory, spin chain models and so on. We call this type of solutions type-I.

Based on Ref. [21] there appears a new type of solutions called type-II [22–25]. By introducing a new variable θ as $\cos\theta = \frac{1+x}{\sqrt{2(1+x^2)}}$ and $\sin\theta = \frac{1-x}{\sqrt{2(1+x^2)}}$, we have

$$\breve{R}_i(\theta) = e^{\theta\gamma_i\gamma_{i+1}} = \cos\theta + \sin\theta\gamma_i\gamma_{i+1}, \tag{9}$$

then the YBE reads [26]:

$$\breve{R}_i(\theta_1)\breve{R}_{i+1}(\theta_2)\breve{R}_i(\theta_3) = \breve{R}_{i+1}(\theta_3)\breve{R}_i(\theta_2)\breve{R}_{i+1}(\theta_1), \tag{10}$$

with the constraint for three parameters θ_1, θ_2 and θ_3 :

$$\tan\theta_2 = \frac{\tan\theta_1 + \tan\theta_3}{1 + \tan\theta_1\tan\theta_3}, \tag{11}$$

i.e. the Lorentzian additivity by $\tan\theta = \frac{1}{c}u$. It is well known that the physical meaning of θ is to describe entangling degree, which is $|\sin 2\theta|$ for 2-qubit [23]. The type-II solution of YBE $\breve{R}_i(\theta)$ means the operation between two Majorana fermions, γ_i and γ_{i+1}. Because γ_i's satisfy Clifford algebraic relations:

$$\{\gamma_i, \gamma_j\} = 2\delta_{ij}. \tag{12}$$

Then the solution $\breve{R}_i(\theta) = e^{\theta\gamma_i\gamma_{i+1}}$ transforms the Majorana fermions γ_i and γ_{i+1} in the following way:

$$\breve{R}_i(\theta)\gamma_i\breve{R}_i^\dagger(\theta) = \cos 2\theta\gamma_i - \sin 2\theta\gamma_{i+1}, \tag{13}$$
$$\breve{R}_i(\theta)\gamma_{i+1}\breve{R}_i^\dagger(\theta) = \sin 2\theta\gamma_i + \cos 2\theta\gamma_{i+1}. \tag{14}$$

Since the solution of Yang-Baxter equation can be expressed in Majorana form, the following problems arise: (i) How to understand the Γ-operator intuitively on the basis of the concrete MF model generated by YBE; (ii) How to obtain the 3-body Hamiltonian, which possesses the doubling degeneracy, from YBE; (iii) What is the relationship between Γ-operator (as well as extended Γ') and the solution $\breve{R}_i(\theta)$ of YBE.

In this paper, we show that the emergent Majorana operator Γ is a new symmetry of $\breve{R}(\theta)$ as well as Yang-Baxter equation. Due to the symmetry, the 3-body Hamiltonian derived from YBE holds Majorana doubling. We also present a new realization of doubling degeneracy for Majorana mode. Moreover, we discuss the topological phase in the "superconducting" chain. The generation of Greenberger-Horne-Zeilinger (GHZ) state via the approach of YBE is also discussed.

RESULTS

Topological phase in the derived "superconducting" chain. The topological phase transition in the derived "superconducting" chain based on YBE is discussed. We find that our chain model is exactly the same as 1D Kitaev model. Let us first give a brief introduction to 1D Kitaev model.

1D Kitaev's toy model is one of the simplest but the most representative model for Majorana mode [1, 4]. The model is a quantum wire with N sites lying on the surface of three dimensional p-wave superconductor, and each site is either empty or occupied by an electron with a fixed spin direction. Then the Hamiltonian is expressed as the following form:

$$\begin{aligned}\hat{H}_k = \sum_j^N \quad & [-\mu\left(a_j^\dagger a_j - \tfrac{1}{2}\right) - \omega\left(a_j^\dagger a_{j+1} + a_{j+1}^\dagger a_j\right) \\ & + \Delta a_j a_{j+1} + \Delta^* a_{j+1}^\dagger a_j^\dagger]. \end{aligned} \tag{15}$$

Here $a_j^\dagger$, a_j represent spinless ordinary fermion, ω is hopping amplitude, μ is chemical potential, and $\Delta = |\Delta|e^{-i\varphi}$ is induced superconducting gap. Define Majorana fermion operators:

$$\gamma_{2j-1} = e^{i\frac{\varphi}{2}} a_j^\dagger + e^{-i\frac{\varphi}{2}} a_j, \tag{16}$$

$$\gamma_{2j} = ie^{i\frac{\varphi}{2}} a_j^\dagger - ie^{-i\frac{\varphi}{2}} a_j, \tag{17}$$

which satisfy the relations:

$$\gamma_m^\dagger = \gamma_m, \quad \{\gamma_l, \gamma_m\} = 2\delta_{lm}, \quad l, m = 1, \ldots 2N. \tag{18}$$

Then the Hamiltonian is transformed into the Majorana form:

$$\begin{aligned}\hat{H}_k = \tfrac{\mathrm{i}}{2} \sum_j \quad & [-\mu\gamma_{2j-1}\gamma_{2j} + (\omega + |\Delta|)\,\gamma_{2j}\gamma_{2j+1} \\ & + (-\omega + |\Delta|)\,\gamma_{2j-1}\gamma_{2j+2}]. \end{aligned} \tag{19}$$

An interesting case is $\mu = 0$, $\omega = |\Delta|$. In this case, the Hamiltonian turns into Majorana mode corresponding to topological phase:

$$\hat{H}_k = \mathrm{i}\omega \sum_j \gamma_{2j}\gamma_{2j+1}. \tag{20}$$

The above Hamiltonian has two degenerate ground states, $|0\rangle$ and $|1\rangle = d^\dagger|0\rangle$. Here $d^\dagger = e^{-\mathrm{i}\varphi/2}(\gamma_1 - \mathrm{i}\gamma_{2N})/2$ is a non-local ordinary fermion. The degenerate states can be used for topological quantum memory qubits that are immune to local errors.

Now let us construct the "superconducting" chain based on the solution $\breve{R}_i(\theta)$ of YBE. We imagine that a unitary evolution is governed by $\breve{R}_i(\theta)$. If only θ in unitary operator $\breve{R}_i(\theta)$ is time-dependent, we can express a state $|\psi(t)\rangle$ as $|\psi(t)\rangle = \breve{R}_i|\psi(0)\rangle$. Taking the Schrödinger equation $i\hbar\frac{\partial}{\partial t}|\psi(t)\rangle = \hat{H}(t)|\psi(t)\rangle$ into account, one obtains:

$$i\hbar\tfrac{\partial}{\partial t}[\breve{R}_i|\psi(0)\rangle] = \hat{H}(t)\breve{R}_i|\psi(0)\rangle. \tag{21}$$

Then the Hamiltonian $\hat{H}_i(t)$ related to the unitary operator $\breve{R}_i(\theta)$ is obtained:

$$\hat{H}_i(t) = \mathrm{i}\hbar\tfrac{\partial \breve{R}_i}{\partial t}\breve{R}_i^{-1}. \tag{22}$$

Substituting $\breve{R}_i(\theta) = \exp(\theta\gamma_i\gamma_{i+1})$ into equation (22), we have:

$$\hat{H}_i(t) = \mathrm{i}\hbar\dot{\theta}\gamma_i\gamma_{i+1}. \tag{23}$$

This Hamiltonian describes the interaction between i-th and $(i+1)$-th sites with the parameter $\dot{\theta}$. Indeed, when $\theta = n \times \frac{\pi}{4}$, the unitary evolution corresponds to the braiding progress of two nearest Majorana fermion sites in the system, here n is an integer and signifies the times of braiding operation.

If we only consider the nearest-neighbour interactions between MFs and extend equation (23) to an inhomogeneous chain with 2N sites, the derived "superconducting" chain model is expressed as:

$$\hat{H} = \mathrm{i}\hbar\sum_{k=1}^{N}(\dot{\theta}_1\gamma_{2k-1}\gamma_{2k} + \dot{\theta}_2\gamma_{2k}\gamma_{2k+1}), \tag{24}$$

with $\dot{\theta}_1$ and $\dot{\theta}_2$ describing odd-even and even-odd pairs, respectively.

Now we give a brief discussion about the above chain model in two cases (see Fig.1):

1. $\dot{\theta}_1 > 0, \dot{\theta}_2 = 0$.

   In this case, the Hamiltonian is:

$$\hat{H}_1 = \mathrm{i}\hbar \sum_k^N \dot{\theta}_1 \gamma_{2k-1}\gamma_{2k}. \tag{25}$$

   As defined in equation (16) and (17), the Majorana operators γ_{2k-1} and γ_{2k} come from the same ordinary fermion site k, $\mathrm{i}\gamma_{2k-1}\gamma_{2k} = 2a_k^\dagger a_k - 1$ ($a_k^\dagger$ and a_k are spinless ordinary fermion operators). $\hat{H}_1$ simply means the total occupancy of ordinary fermions in the chain and has U(1) symmetry, $a_j \to e^{i\phi}a_j$. Specifically, when $\theta_1(t) = \frac{\pi}{4}$, the unitary evolution $e^{\theta_1\gamma_{2k-1}\gamma_{2k}}$ corresponds to the braiding operation of two Majorana sites from the same k-th ordinary fermion site. The ground state represents the ordinary fermion occupation number 0. In comparison to 1D Kitaev model, this Hamiltonian corresponds to the trivial case of Kitaev's. In Fig.1, this Hamiltonian is described by the intersecting lines above the dashed line, where the intersecting lines correspond to interactions. The unitary evolution of the system $e^{-i\int \hat{H}_1 dt}$ stands for the exchange process of odd-even Majorana sites.

2. $\dot{\theta}_1 = 0, \dot{\theta}_2 > 0$.

   In this case, the Hamiltonian is:

$$\hat{H}_2 = \mathrm{i}\hbar \sum_k^N \dot{\theta}_2 \gamma_{2k}\gamma_{2k+1}. \tag{26}$$

   This Hamiltonian corresponds to the topological phase of 1D Kitaev model and has $\mathbb{Z}_2$ symmetry, $a_j \to -a_j$. Here the operators γ_1 and γ_{2N} are absent in $\hat{H}_2$, which is illustrated by the crossing under the dashed line in Fig.1. The Hamiltonian has two degenerate ground state, $|0\rangle$ and $|1\rangle = d^\dagger|0\rangle$, $d^\dagger = e^{-i\varphi/2}(\gamma_1 - i\gamma_{2N})/2$. This mode is the so-called Majorana mode in 1D Kitaev chain model. When $\theta_2(t) = \frac{\pi}{4}$, the unitary evolution $e^{\theta_2\gamma_{2k}\gamma_{2k+1}}$ corresponds to the braiding operation of two Majorana sites γ_{2k} and γ_{2k+1} from k-th and $(k+1)$-th ordinary fermion sites, respectively.

Thus we conclude that our Hamiltonian derived from $\breve{R}_i(\theta(t))$ corresponding to the braiding of nearest Majorana fermion sites is exactly the same as the 1D wire proposed by Kitaev, and $\dot{\theta}_1 = \dot{\theta}_2$ corresponds to the phase transition point in the "superconducting" chain. By choosing different time-dependent parameter θ_1 and θ_2, we find that the Hamiltonian $\hat{H}$ corresponds to different phases.

New realization of Majorana Doubling based on Γ-operator. The important progress had been made to establish the complete algebra for the Majorana doubling by introducing the emergent Majorana operator Γ [8]:

$$\Gamma = -\mathrm{i}\gamma_1\gamma_2\gamma_3. \tag{27}$$

In Ref. [8], the concreted realization of the operators was presented in terms of Pauli matrices. On the other hand, as pointed out in Ref. [27], there is the transformation between the natural basis and Bell basis for

$$|\Phi_0\rangle = (|\downarrow\downarrow\rangle, |\uparrow\downarrow\rangle, |\downarrow\uparrow\rangle, |\uparrow\uparrow\rangle)^T, \tag{28}$$
$$|\Psi\rangle = (|\Psi_+\rangle, |\Phi_+\rangle, |\Phi_-\rangle, |\Psi_-\rangle)^T, \tag{29}$$

where

$$\big|\Psi_+\big\rangle = \frac{1}{\sqrt{2}}\big(\big|\uparrow\uparrow\big\rangle + \big|\downarrow\downarrow\big\rangle\big), \qquad \big|\Phi_+\big\rangle = \tfrac{1}{\sqrt{2}}\big(\big|\uparrow\downarrow\big\rangle + \big|\uparrow\downarrow\big\rangle\big), \tag{30}$$
$$\big|\Psi_-\big\rangle = \frac{1}{\sqrt{2}}\big(\big|\downarrow\uparrow\big\rangle - \big|\uparrow\downarrow\big\rangle\big), \qquad \big|\Phi_-\big\rangle = \tfrac{1}{\sqrt{2}}\big(\big|\uparrow\uparrow\big\rangle - \big|\downarrow\downarrow\big\rangle\big) \tag{31}$$

through the matrix B_{II}:

$$\big|\Psi\big\rangle = B_{II}\big|\Phi_0\big\rangle, \tag{32}$$

where

$$B_{II}=\frac{1}{\sqrt{2}}\begin{bmatrix}1&0&0&1\\0&1&1&0\\0&-1&1&0\\-1&0&0&1\end{bmatrix}=\frac{1}{\sqrt{2}}(I+M)\quad(M^2=-1) \tag{33}$$

and

$$M_iM_{i\pm1} = -M_{i\pm1}M_i,\quad M^2=-I, \tag{34}$$
$$M_iM_j = M_jM_{i,}\quad |i-j|\geq 2 \tag{35}$$

which forms "extra special 2-group". Obviously, M is extension of i for $\mathrm{i}^2=-1$.

An interesting observation is [28]:

$$M=-\mathrm{i}\hat{C} \tag{36}$$

where $\hat{C}$ is the charge conjugate operator in Majorana spinor. The eigenstates of $\hat{C}$ take the forms

$$\hat{C}|\xi_\pm\rangle=\mp|\xi_\pm\rangle,\qquad \hat{C}|\eta_\pm\rangle=\mp|\eta_\pm\rangle, \tag{37}$$

where

$$|\xi_\pm\rangle=\frac{1}{\sqrt{2}}(|\uparrow\uparrow\rangle\pm\mathrm{i}|\downarrow\downarrow\rangle), \tag{38}$$
$$|\eta_\pm\rangle=\frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle\pm\mathrm{i}|\downarrow\uparrow\rangle). \tag{39}$$

Here we would like to give an intuitive interpretation of the operator Γ in Ref. [8] by taking a new set of $D_i\,(i=1,2,3)$ in stead of γ_i, and show how it gives rise to the Majorona doubling with explicit realization.

We follow the concrete realization for γ_j given in Ref. [8], (in this paper I is 2×2 identity matrix)

$$\gamma_1=\sigma_1\otimes I,\ \gamma_2=\sigma_3\otimes I,\ \gamma_3=\sigma_2\otimes\sigma_1, \tag{40}$$
$$P=\sigma_2\otimes\sigma_3, \tag{41}$$
$$\Gamma=-\mathrm{i}\gamma_1\gamma_2\gamma_3=-I\otimes\sigma_1. \tag{42}$$

In our notation, $\gamma_3=-\hat{C}$, i.e. (38) and (39) are eigenstates of γ_3. It is easy to find

$$\gamma_1|\xi_\pm\rangle=\pm\mathrm{i}|\eta_\mp\rangle,\quad \gamma_1|\eta_\pm\rangle=\pm\mathrm{i}|\xi_\mp\rangle; \tag{43}$$
$$\gamma_2|\xi_\pm\rangle=|\xi_\mp\rangle,\qquad \gamma_2|\eta_\pm\rangle=|\eta_\mp\rangle; \tag{44}$$
$$\gamma_3|\xi_\pm\rangle=\pm|\xi_\pm\rangle,\quad \gamma_3|\eta_\pm\rangle=\pm|\eta_\pm\rangle; \tag{45}$$
$$P|\xi_\pm\rangle=\mp|\eta_\mp\rangle,\quad P|\eta_\pm\rangle=\pm|\xi_\mp\rangle; \tag{46}$$
$$\Gamma|\xi_\pm\rangle=-|\eta_\pm\rangle,\qquad \Gamma|\eta_\pm\rangle=-|\xi_\pm\rangle. \tag{47}$$

In the derivation of (43)-(47), the relations $\sigma_1=(S^+ + S^-)$ and $\sigma_2=\frac{1}{\mathrm{i}}(S^+-S^-)$ have been used where $S^\pm=S_1\pm\mathrm{i}S_2$. To show the importance of Γ-operator we define new Clifford algebra $\{D_i,D_j\}=2\delta_{ij}$, where $D_1=\gamma_2,\ D_2=\Gamma\gamma_1,\ D_3=\gamma_3$. It is interesting to find that

$$D_j|\xi\rangle = \sigma_j|\xi\rangle,\quad D_j|\eta\rangle=\sigma_j|\eta\rangle,\qquad (j=1,2,3) \tag{48}$$
$$|\xi\rangle = \begin{pmatrix}|\xi_+\rangle\\|\xi_-\rangle\end{pmatrix},\quad |\eta\rangle=\begin{pmatrix}|\eta_+\rangle\\|\eta_-\rangle\end{pmatrix}. \tag{49}$$

Namely, by acting D_j on$|\xi\rangle$ or $|\eta\rangle$, the representation is exactly Pauli matrices, i.e. belonging to SU(2) algebra. It can be checked that

$$D_1D_2=-\mathrm{i}\Sigma_2,\quad D_2D_3=-\mathrm{i}\Sigma_3,\quad D_1D_3=-\mathrm{i}\Sigma_1, \tag{50}$$

where Σ_i form the reducible representation of $SU(2)$:

$$\Sigma_1 = \sigma_1 \otimes \sigma_1, \quad \Sigma_2 = \sigma_2 \otimes \sigma_1, \quad \Sigma_3 = \sigma_3 \otimes I. \tag{51}$$

The introduced interacting Hamiltonian $H_B = -\mathrm{i}(\alpha D_1 D_2 + \beta D_2 D_3 + \kappa D_3 D_1)$ can be recast to

$$H_B = -(\alpha_1 \Sigma_1 + \alpha_2 \Sigma_2 + \alpha_3 \Sigma_3), \tag{52}$$

where $\alpha_1 = -\kappa,\ \alpha_2 = \alpha,\ \alpha_3 = \beta$. Noting that $D_1 D_2 D_3 = -\mathrm{i} I \otimes I$, i.e. trivial. The direct check gives:

$$\Big[\Gamma, \Sigma_j\Big] = 0, \qquad (j = 1, 2, 3) \tag{53}$$

and

$$\Big[\Sigma_j, \Sigma_k\Big] = \mathrm{i}\epsilon_{jkl}\Sigma_l.$$

Then the H_B can be written in the form:

$$\begin{aligned} H_B &= E\overrightarrow{n}\cdot\overrightarrow{\Sigma}, \qquad (\overrightarrow{\Sigma}^2 = I), && (54)\\ \overrightarrow{n} &= (\sin\zeta\cos\varphi,\ \sin\zeta\sin\varphi,\ \cos\zeta), && (55)\\ \cos\zeta &= -\alpha_3/E, \quad \tan\varphi = \alpha_2/\alpha_1. && (56) \end{aligned}$$

Obviously, $\overrightarrow{\Sigma}$ is reducible 4-d representation of SU(2). Explicitly,

$$\begin{aligned} \overrightarrow{n}\cdot\overrightarrow{\Sigma} &= M_1 + M_2 \\ &= \begin{bmatrix} \cos\zeta & 0 & 0 & \sin\zeta e^{-i\varphi} \\ 0 & \cos\zeta & \sin\zeta e^{-i\varphi} & 0 \\ 0 & \sin\zeta e^{i\varphi} & -\cos\zeta & 0 \\ \sin\zeta e^{i\varphi} & 0 & 0 & -\cos\zeta \end{bmatrix}, \end{aligned} \tag{57}$$

where

$$M_1 = \begin{bmatrix} \cos\zeta & 0 & 0 & \sin\zeta e^{-i\varphi} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \sin\zeta e^{i\varphi} & 0 & 0 & -\cos\zeta \end{bmatrix}, \tag{58}$$

$$M_2 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & \cos\zeta & \sin\zeta e^{-i\varphi} & 0 \\ 0 & \sin\zeta e^{i\varphi} & -\cos\zeta & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}. \tag{59}$$

Rewriting M_1 and M_2 in the form of Pauli matrices, we have

$$M_1 = \cos\zeta\frac{\sigma_3 \otimes I + I \otimes \sigma_3}{2} + \sin\zeta(e^{-i\varphi}\sigma^+ \otimes \sigma^+ + e^{i\varphi}\sigma^- \otimes \sigma^-), \tag{60}$$

$$M_2 = \cos\zeta\frac{\sigma_3 \otimes I - I \otimes \sigma_3}{2} + \sin\zeta(e^{-i\varphi}\sigma^+ \otimes \sigma^- + e^{i\varphi}\sigma^- \otimes \sigma^+). \tag{61}$$

Now the meaning of H_B is manifest: 4-dimension is quite different from 2-dimension. The "edge block" leads to M_1 with superconducting type of Hamiltonian whereas "interior block" M_2 is connected with the usual spin chain. It is easy to find the eigenstates of M_1 and M_2:

$$M_1\Big|\psi_1\Big\rangle = \Big|\psi_1\Big\rangle, \qquad M_2\Big|\psi_2\Big\rangle = \Big|\psi_2\Big\rangle, \tag{62}$$

where

$$\Big|\psi_1\Big\rangle = \begin{bmatrix} \cos\frac{\zeta}{2} \\ 0 \\ 0 \\ \sin\frac{\zeta}{2}\, e^{i\varphi} \end{bmatrix}, \qquad \Big|\psi_2\Big\rangle = \begin{bmatrix} 0 \\ \cos\frac{\zeta}{2} \\ \sin\frac{\zeta}{2}\, e^{i\varphi} \\ 0 \end{bmatrix}. \tag{63}$$

Acting Γ on (63) it yields

$$\Gamma\Big|\psi_1\Big\rangle = -\Big|\psi_2\Big\rangle, \qquad \Gamma\Big|\psi_2\Big\rangle = -\Big|\psi_1\Big\rangle. \tag{64}$$

So Γ transforms between $\big|\psi_1\big\rangle$ and $\big|\psi_2\big\rangle$ that holds for the same energy. It never occurs in 2 dimensions. Meanwhile, equation (53) shows that Γ commutes with the Hamiltonian H_B, which means that Γ-transformation does not change the property of Hamiltonian H_B. This example shows that operator Γ is crucial in leading to Majorana doubling in dimensions ≥ 4. With the new defination of D_2, we should define a new parity operator:

$$P_B = \sigma_3 \otimes \sigma_2. \tag{65}$$

Direct check gives the complete set of algebra

$$\{D_i, D_j\} = 0, \tag{66}$$
$$\Gamma^2 = I,\ [\Gamma, D_j] = 0,\ [\Gamma, H_B] = 0, \tag{67}$$
$$P_B^2 = I,\ [P_B, D_j] = 0, \tag{68}$$
$$\{\Gamma, P_B\} = 0,\ [P_B, H_B] = 0, \tag{69}$$
$$[\Gamma, \Sigma_j] = 0,\ [\Sigma_j, \Sigma_k] = \mathrm{i}\epsilon_{jkl}\Sigma_l,\ (j,k,l = 1,2,3)\,. \tag{70}$$

It is noteworthy that the introduced P_B in equation (68) commutes with D_j instead of the anti commuting relation between P and γ_j. And P_B still anticommutes with Γ. Acting P_B on the eigenstates $|\psi_1\rangle$ and $|\psi_2\rangle$, it follows

$$P_B\,|\psi_1\rangle = \mathrm{i}\,|\psi_2\rangle\,, \quad P_B\,|\psi_2\rangle = -\mathrm{i}\,|\psi_1\rangle\,. \tag{71}$$

In such a concrete realization Γ plays the essential role. The Hamiltonian (54) formed by (52) looks a typical nuclear resonant model in 4 dimensions. Only the higher dimensions allow the operator Γ leading to the doubling degeneracy.

Majorana doubling in 3-body Hamiltonian based on YBE. Now we discuss the interaction of 3 Majorana fermions based on YBE.

It is well known that $\breve{R}_i(\theta)$ describes the 2-body interaction. And the physical meaning of Yang-Baxter equation is that the interaction of the three bodies can be decomposed into three 2-body interactions:

$$\begin{aligned}\breve{R}_{123}(\theta_1,\theta_2,\theta_3) &= \breve{R}_{12}(\theta_1)\breve{R}_{23}(\theta_2)\breve{R}_{12}(\theta_3)\\ &= \breve{R}_{23}(\theta_3)\breve{R}_{12}(\theta_2)\breve{R}_{23}(\theta_1).\end{aligned}$$

Because of the constraint in equation (11), $\breve{R}_{123}$ depends only on two free parameters and has the following form [29]:

$$\breve{R}_{123}(\eta,\beta) = e^{\eta\left(\vec{n}\cdot\vec{\Lambda}\right)}, \tag{72}$$

where

$$\begin{aligned}\cos\eta &= \cos\theta_2\cos\left(\theta_1+\theta_3\right),\\ \sin\eta &= \sin\theta_2\sqrt{1+\cos^2(\theta_1-\theta_3)},\\ \vec{n} &= \Big(\ \tfrac{1}{\sqrt{2}}\cos\beta,\ \ \tfrac{1}{\sqrt{2}}\cos\beta,\ \ \sin\beta\ \Big),\\ \vec{\Lambda} &= \big(\ \gamma_1\gamma_2,\ \ \gamma_2\gamma_3,\ \ \gamma_1\gamma_3\ \big),\\ \cos\beta &= \tfrac{\sqrt{2}\cos(\theta_1-\theta_3)}{\sqrt{1+\cos^2(\theta_1-\theta_3)}},\\ \sin\beta &= \tfrac{-\sin(\theta_1-\theta_3)}{\sqrt{1+\cos^2(\theta_1-\theta_3)}}.\end{aligned}$$

Here the parameters θ_1 and θ_3 are replaced by η and β. $\breve{R}_{123}(\eta,\beta)$ is also a unitary operator and describes the interaction of three Majorana operators.

We suppose that the parameter η is time-dependent and β is time-independent in $\breve{R}_{123}(\eta,\beta)$, then the desired 3-body Hamiltonian can be obtained from equation (22):

$$\begin{aligned}\hat{H}_{123}(t) &= \mathrm{i}\hbar\tfrac{\partial\breve{R}_{123}}{\partial t}\breve{R}_{123}^{-1}\\ &= \mathrm{i}\hbar\dot{\eta}\left[\tfrac{1}{\sqrt{2}}\cos\beta(\gamma_1\gamma_2+\gamma_2\gamma_3)+\sin\beta\gamma_1\gamma_3\right].\end{aligned} \tag{73}$$

The constructed Hamiltonian, which has been mentioned in Ref. [7, 8], describes the 2-body interactions among the three Majorana operators. It describes the effective interaction in a T-junction formed by three quantum wires. In Ref. [8], it has been shown that the above Hamiltonian, which commutes with emergent Majorana operator $\Gamma = -\mathrm{i}\gamma_1\gamma_2\gamma_3$, holds Majorana doubling. From the viewpoint of YBE, the intrinsic commutation relation is between Γ and the solution of YBE $\breve{R}_i(\theta) = e^{\theta\gamma_i\gamma_{i+1}}$. It is shown that:

$$[\Gamma, \breve{R}_i(\theta)] = 0, \quad (i = 1, 2). \tag{74}$$

Indeed, the above commutation relation indicates that emergent Majorana operator Γ is a new symmetry of the solution $\breve{R}_i(\theta)$ of YBE. It is due to the decomposition of 3-body interaction into three 2-body interactions via the approach of YBE that the derived Hamiltonian holds Majorana doubling.

The extended emergent Majorana mode Γ' supporting odd number N of Majorana operators [8] is,

$$\Gamma' \equiv \mathrm{i}^{N(N-1)/2} \prod_{j=1}^{N} \gamma_j. \tag{75}$$

It is easy to check that:

$$[\Gamma', B_i] = 0, \quad (i = 1, 2, ...N-1), \tag{76}$$

where $B_i = e^{\frac{\pi}{4}\gamma_i\gamma_{i+1}}$ is the generator of the braid group B_N. The commutation relation indicates that Γ' plays the role of an invariant in the braid group B_N.

Generation of 2n-qubit GHZ state via YBE. Quantum entanglement plays an important role in quantum information theory and has been discussed in both theoretical [30] and experimental [31, 32] aspects for a long time. There are various ways in describing different types of entanglement. It is also well known that the relationship between Yang-Baxter equation and 2-qubit entangled state as well as 3-qubit entanglement has been discussed in Ref. [22, 23, 29, 34]. Here we construct high dimensional matrix representation of solution to Yang-Baxter equation and discuss how it generates $2N$-qubit GHZ state for odd N. In previous section, we present Clifford algebric relation for different Majorana operators,

$$\{\gamma_i, \gamma_j\} = 2\delta_{ij}. \tag{77}$$

It can be used for constructing solution to YBE:

$$\breve{R}_i(\theta) = \exp(\theta\gamma_i\gamma_{i+1}). \tag{78}$$

The representation of γ_i in the Majorana form is given by:

$$\gamma_{2j-1} = e^{i\varphi}a_j^\dagger + e^{-i\varphi}a_j, \tag{79}$$

$$\gamma_{2j} = ie^{i\varphi}a_j^\dagger - ie^{-i\varphi}a_j. \tag{80}$$

Then by constructing Yang-Baxter chain, we find its similarity to 1D Kitaev model.

Indeed, the 4D-matrix representation is equivalent to the Majorana fermion representation under Jordan-Wigner transformation. In other words, we can express γ_i by matrix directly. For three operators γ_1, γ_2 and γ_3 satisfying Clifford algebra, its 4D matrix representation has been presented in Ref. [8]:

$$\begin{aligned}
\gamma_1 &= \sigma_1 \otimes I, \\
\gamma_2 &= \sigma_3 \otimes I, \\
\gamma_3 &= \sigma_2 \otimes \sigma_1,
\end{aligned}$$

here σ_i are Pauli matrices.

What we are interested in is constructing higher dimensional matrix representation of γ_i. Taking 8D representation as an example, γ_i is:

$$\begin{aligned}
\gamma_1 &= \sigma_1 \otimes I \otimes I, \\
\gamma_2 &= \sigma_3 \otimes \sigma_1 \otimes I, \\
\gamma_3 &= \sigma_3 \otimes \sigma_3 \otimes \sigma_1.
\end{aligned}$$

Then the matrix form of emergent Majorana mode Γ [8] is,

$$\Gamma = -\mathrm{i}\gamma_1\gamma_2\gamma_3 = -\sigma_1 \otimes \sigma_2 \otimes \sigma_1. \tag{81}$$

The Hamiltonian supporting three Majorana operators has been defined in equation (2):

$$\begin{aligned} H_{\text{int}} &= -\mathrm{i}\left(\alpha\gamma_1\gamma_2 + \beta\gamma_2\gamma_3 + \kappa\gamma_3\gamma_1\right) \\ &= -\alpha\sigma_2 \otimes \sigma_1 \otimes I - \beta I \otimes \sigma_2 \otimes \sigma_1 + \kappa\sigma_2 \otimes \sigma_3 \otimes \sigma_1. \end{aligned} \tag{82}$$

Obviously, Γ commutes with the Hamiltonian H_{int}.

Let us extend Γ to N sites Γ_i, which should also satisfy Clifford algebra $\{\Gamma_i, \Gamma_j\} = 2\delta_{ij}$. The Γ_i has the following form:

$$\begin{aligned} \gamma_{3i-2} &= (\sigma_3 \otimes \sigma_3 \otimes \sigma_3)^{\otimes(i-1)} \otimes \sigma_1 \otimes I \otimes I \cdots, \\ \gamma_{3i-1} &= (\sigma_3 \otimes \sigma_3 \otimes \sigma_3)^{\otimes(i-1)} \otimes \sigma_3 \otimes \sigma_1 \otimes I \cdots, \\ \gamma_{3i} &= (\sigma_3 \otimes \sigma_3 \otimes \sigma_3)^{\otimes(i-1)} \otimes \sigma_3 \otimes \sigma_3 \otimes \sigma_1 \cdots, \end{aligned}$$

$$\begin{aligned} \Gamma_i &= -\mathrm{i}\gamma_{3i-2}\gamma_{3i-1}\gamma_{3i} \\ &= -(\sigma_3 \otimes \sigma_3 \otimes \sigma_3)^{\otimes(i-1)} \otimes \sigma_1 \otimes \sigma_2 \otimes \sigma_1 \otimes I \cdots. \end{aligned} \tag{83}$$

Then we have:

$$\Gamma_i\Gamma_{i+1} = -\mathrm{i}I^{\otimes 3(i-1)} \otimes (\sigma_2 \otimes \sigma_1)^{\otimes 3} \otimes I \otimes I \cdots. \tag{84}$$

It is easy to check that $e^{\theta\Gamma_i\Gamma_{i+1}}$ is the 4^3-D matrix solution of YBE, we denote it by $\breve{R}_i^3(\theta)$,

$$\breve{R}_i^3(\theta) = \cos\theta I^{\otimes 6} - \mathrm{i}\sin\theta(\sigma_2 \otimes \sigma_1)^{\otimes 3}. \tag{85}$$

By acting $\breve{R}_i^3(\theta)$ on 6-qubit natural basis, such as $|\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow\rangle$, we have:

$$\breve{R}_i^3(\theta)|\uparrow\rangle^{\otimes 6} = \cos\theta|\uparrow\rangle^{\otimes 6} - \sin\theta|\downarrow\rangle^{\otimes 6}. \tag{86}$$

This state represents a type of 6-qubit entangled states. In the case of $\theta = \frac{\pi}{4}$, the generated state is 6-qubit GHZ state, and $\breve{R}_i^3(\theta = \frac{\pi}{4}) = e^{\frac{\pi}{4}\Gamma_i\Gamma_{i+1}}$ can be regarded as one braiding operation of two emergent Majorana operator Γ_i and Γ_{i+1}.

Now we generalize the 4^3-D matrix solution of YBE to 4^n with n odd. The extended Majorana operator supporting any odd number n of Majorana operators reads,

$$\Gamma^n = \Gamma' \equiv \mathrm{i}^{n(n-1)/2}\prod_{j=1}^{n}\gamma_j, \tag{87}$$

where the constraint of Clifford algebra $\{\Gamma_i^n, \Gamma_j^n\} = 2\delta_{ij}$ leads to the odd number n. Γ_i^n can be expressed as:

$$\Gamma_i^n = -(\sigma_3)^{\otimes n(i-1)} \otimes (\sigma_1 \otimes \sigma_2)^{\otimes\frac{n-1}{2}} \otimes \sigma_1 \otimes I \otimes I \cdots. \tag{88}$$

Then we have

$$\Gamma_i^n\Gamma_{i+1}^n = -(\mathrm{i})I^{\otimes n(i-1)} \otimes (\sigma_2 \otimes \sigma_1)^{\otimes n} \otimes I \otimes I \cdots. \tag{89}$$

The 4^n-D (n odd) matrix representation of solution to YBE is:

$$\begin{aligned} \breve{R}_i^n(\theta) &= e^{\theta\Gamma_i^n\Gamma_{i+1}^n} \\ &= \cos\theta I^{\otimes 2n} - \mathrm{i}\sin\theta(\sigma_2 \otimes \sigma_1)^{\otimes n} \quad (\text{n odd}). \end{aligned} \tag{90}$$

Consequently, we generate the following state by acting $\breve{R}_i^n(\theta)$ on the $2n$(n odd)-qubit natural state $|\uparrow\rangle^{\otimes 2n}$:

$$\breve{R}_i^n(\theta)|\uparrow\rangle^{\otimes 2n} = \cos\theta|\uparrow\rangle^{\otimes 2n} - \sin\theta|\downarrow\rangle^{\otimes 2n}. \tag{91}$$

When $\theta = \frac{\pi}{4}$, the generated state turns into $2n$-qubit GHZ state for odd n.

DISCUSSION

In this paper, based on the solution of YBE in Majorana form, we discuss the topological phase transition in the derived "superconducting" chain and the Majorana doubling in 3-body Hamiltonian as well as the generation of 2n-qubit GHZ-type entangled states. Unlike the braid operator, the solution $\breve{R}_i(\theta)$ of YBE is parameter-dependent. Hence the unitary operator $\breve{R}_i(\theta)$ can be used for generating the "superconducting" chain and the Majorana doubling in 3-body Hamiltonian. Indeed, the derived chain(25,26) describes the braiding transformation of nearest-neighbour Majorana sites for $\theta_1 = \frac{\pi}{4}$ (or $\theta_2 = \frac{\pi}{4}$). We also find that the 3-body Hamiltonian $\hat{H}_{123}$ derived from $\breve{R}_{123}$ holds Majorana doubling. From the viewpoint of YBE, the commutation relation $[\Gamma, \hat{H}_{123}] = 0$ can be explained by $[\Gamma, \breve{R}_i(\theta)] = 0$ (i=1,2), where $\breve{R}_i(\theta)$ is the solution of YBE. In other words, it is the Γ-symmetry of $\breve{R}(\theta)$ that leads to the Γ-symmetry of $\hat{H}_{123}$. The commutation relation can also be generalized to the extended Γ'-operator(87) for odd N sites, $[\Gamma', B_i] = 0$ ($i = 1, 2, ...N-1$), hence Γ' is an invariant of the braid group B_N.

We present a new realization of Majorana doubling based on emergent Majorana mode and show the role of Γ in leading to the doubling degeneracy of H_B intuitively. We also make use of the extended Γ'-operator to construct high dimensional matrix representation of solution to YBE. By acting the high dimensional matrix representation of solution of YBE on natural basis, we generate the GHZ-type entangled state. Thus we conclude that the braiding process of the extended Γ'-operators corresponds to the generation of GHZ entangled state. These results may guide us to find much closer relationship between Yang-Baxter equation and quantum information as well as condensed matter physics.

[1] Kitaev, A.Yu. Unpaired majorana fermions in quantum wires. *Phys. Usp.* **44,** 131 (2001).
[2] Ivanov, D.A. Non-abelian statistics of half-quantum vortices in p-wave superconductors. *Phys. Rev. Lett.* **86,** 268–271 (2001).
[3] Wilczek, F. Majorana returns. *Nature Phys.* **5,** 614–618 (2009).
[4] Leijnse, M. & Flensberg, K. Introduction to topological superconductivity and Majorana fermions. *Semicon. Sci. Tech.* **27,** 124003 (2012).
[5] Kitaev, A. Yu. Fault-tolerant quantum computation by anyons. *Ann. Phys.* **303,** 2–30 (2003).
[6] Kitaev, A. Yu. Anyons in an exactly solved model and beyond. *Ann. Phys.* **321,** 2–111 (2006).
[7] Alicea, J. *et al.* Non-Abelian statistics and topological quantum information processing in 1D wire networks. *Nature Phys.* **7,** 412–417 (2011).
[8] Lee, J. & Wilczek, F. Algebra of majorana doubling. *Phys. Rev. Lett.* **111,** 226402 (2013).
[9] Baxter, R. J. Partition function of the eight-vertex lattice model. *Ann. Phys.* **70,** 193–228 (1972).
[10] Wilczek, F. *Fractional Statistics and Anyon Superconductivity* (World Scientific, 1990).
[11] Yang, C. N. Some exact results for the many-body problem in one dimension with repulsive delta-function interaction. *Phys. Rev. Lett.* **19,** 1312 (1967).
[12] Yang, C. N. S matrix for the one-dimensional N-body problem with repulsive or attractive δ-function interaction. *Phys. Rev.* **168,** 1920 (1968).
[13] Baxter, R. J. *Exactly solved models in statistical mechanics* (Courier Dover Publications, 2007).
[14] Batchelor, M. T. The Bethe ansatz after 75 years. *Phys. Tod.* **60,** 36 (2007).
[15] Ge, M. L. & Yang, C. N.(Eds.) *Braid group, Knot Theory And Statistical Mechanics* (World Scientific, 1989)
[16] Takhtadzhan, L. A. & Faddeev, L. D. The quantum method of the inverse problem and the Heisenberg XYZ model. *Rus. Math. Sur.* **34,** 11–68 (1979).
[17] Faddeev, L. D. Quantum completely integrable models in field theory. *Sov. Sci. Rev. Math. Phys. C* **1,** 107–155 (1981).
[18] Kulish, P. & Sklyanin, E. Quantum spectral transform method recent developments. *Integrable quantum field theories*,61–119(Springer, 1982).
[19] Kulish, P. & Sklyanin, E. K. Solutions of the Yang-Baxter equation. *J. Sov. Math.* **19,** 1596–1620 (1982).
[20] Korepin, V. E. *Quantum inverse scattering method and correlation functions* (Cambridge Univ. Press, 1997).
[21] Hietarinta, J. Solving the two-dimensional constant quantum Yang–Baxter equation. *J. Math. Phys.* **34,** 1725–1756 (1993).
[22] Kauffman, L. H. & Lomonaco, S. J. J. Quantum entanglement and topological entanglement. *New J. Phys.* **4,** 73 (2002).
[23] Chen, J. L., Xue, K., & Ge, M. L. Braiding transformation, entanglement swapping, and Berry phase in entanglement space. *Phys. Rev. A* **7,** 042324 (2007).
[24] Hu, S. W., Xue, K. & Ge, M. L. Optical simulation of the Yang-Baxter equation. *Phys. Rev. A* **78,** 022319 (2008).
[25] Ge, M. L. & Xue, K. Yang-Baxter equations in quantum information. *Int. J. Mod. Phys. B* **26,** 27n28 (2012).
[26] Jimbo, M. *Yang-Baxter Equation in Integrable Systems* (World Scientific, 1990).
[27] Kauffman, L. H. & Lomonaco, S. J. J. Braiding operators are universal quantum gates. *New J. Phys.* **6,** 134 (2004).
[28] Ge, M. L., Yu, L. W., Xue, K. & Zhao, Q. Yang-Baxter equation, Majorana fermions and three body entangling states. *Int. J. Mod. Phys. B* **28,** 1450089 (2014).
[29] Yu, L. W., Zhao, Q. & Ge, M. L. Factorized three-body S-matrix restrained by Yang–Baxter equation and quantum entanglements. *Ann. Phys.* **348,** 106–126 (2014).
[30] Horodecki, R., Horodecki, P., Horodecki, M., & Horodecki, K. Quantum entanglement. *Rev. Mod. Phys.* **81,** 865–942 (2009).
[31] Yamamoto, T., Koashi, M., Ozdemir, S. K., & Imoto, N. Experimental extraction of an entangled photon pair from two identically

decohered pairs. *Nature* **421,** 343–346 (2003).
[32] Lu, C. Y. *et al*. Experimental entanglement of six photons in graph states. *Nature Phys.* **3,** 91–95 (2007).
[33] Yao, X. C. *et al*. Observation of eight-photon entanglement. *Nature Photon.* **6,** 225–228 (2012).
[34] Zhang, Y., Kauffman, L. H. & Ge, M. L. Universal quantum gate, Yang–Baxterization and Hamiltonian. *Int. J. Quan. Inf.* **3,** 669–678 (2005).

Acknowledgments This work is in part supported by NSF of China (Grant No. 11475088).

Author contributions M.L.G. proposed the idea, L.W.Y. performed the calculation and derivation, L.W.Y. and M.L.G. prepared the manuscript, all authors reviewed the manuscript.

Additional information
Competing financial interests: The authors declare no competing financial interests.



Figure 1: **The nearest neighbouring interactions of 2N Majorana sites described by the "superconducting" chain.** Each solid line represents a Majorana site, and the crossing means the interaction. The dashed line divides the interactions into two parts that are described by $\dot{\theta}_1$ and $\dot{\theta}_2$ respectively. When $\dot{\theta}_1 = 0, \dot{\theta}_2 \neq 0$, the first line and the last line are free, and the Hamiltonian corresponds to topological phase.